



Maritime Aerosol Network as
a component of
AERONET

A. Smirnov et al.

This discussion paper is/has been under review for the journal Atmospheric Measurement Techniques (AMT). Please refer to the corresponding final paper in AMT if available.

Maritime Aerosol Network as a component of AERONET – first results and comparison with global aerosol models and satellite retrievals

A. Smirnov^{1,2}, B. N. Holben², D. M. Giles^{1,2}, I. Slutsker^{1,2}, N. T. O'Neill³,
T. F. Eck^{4,2}, A. Macke⁵, P. Croot⁶, Y. Courcoux⁷, S. M. Sakerin⁸, T. J. Smyth⁹,
T. Zielinski¹⁰, G. Zibordi¹¹, J. I. Goes¹², M. J. Harvey¹³, P. K. Quinn¹⁴,
N. B. Nelson¹⁵, V. F. Radionov¹⁶, C. M. Duarte¹⁷, R. Losno¹⁸, J. Sciare¹⁹,
K. J. Voss²⁰, S. Kinne²¹, N. R. Nalli²², E. Joseph²³, K. Krishna Moorthy²⁴,
D. S. Covert²⁵, S. K. Gulev²⁶, G. Milinevsky²⁷, P. Larouche²⁸, S. Belanger²⁹,
E. Horne³⁰, M. Chin³¹, L. A. Remer³², R. A. Kahn³², J. S. Reid³³, M. Schulz¹⁹,
C. L. Heald³⁴, J. Zhang³⁵, K. Lapina³⁴, R. G. Kleidman^{36,32}, J. Griesfeller¹⁹,
B. J. Gaitley³⁷, Q. Tan^{4,2}, and T. L. Diehl^{4,2}

¹Sigma Space Corporation, Lanham, Maryland, USA

²Biospheric Sciences Branch, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



³CARTEL, Université de Sherbrooke, Sherbrooke, Québec, Canada

⁴Goddard Earth Sciences and Technology Center, University of Maryland, Baltimore County, Baltimore, Maryland, USA

⁵Leibniz Institute for Tropospheric Research, Leipzig, Germany

⁶Leibniz Institute of Marine Sciences at the University of Kiel (IFM-GEOMAR), Kiel, Germany

⁷L'Observatoire de Physique de l'Atmosphère de la Réunion (OPAR), Université de la Réunion, Saint Denis de la Réunion, France

⁸Institute of Atmospheric Optics, Russian Academy of Sciences, Siberian Branch, Tomsk, Russia

⁹Plymouth Marine Laboratory, Plymouth, UK

¹⁰Institute of Oceanology, Polish Academy of Sciences, Sopot, Poland

¹¹Institute for Environment and Sustainability, Joint Research Centre, European Commission, Ispra, Italy

¹²Bigelow Laboratory for Ocean Sciences, West Boothbay Harbor, Maine, USA

¹³National Institute of Water and Atmospheric Research, Wellington, New Zealand

¹⁴NOAA Pacific Marine Environmental Laboratory, Seattle, Washington, USA

¹⁵Institute for Computational Earth System Science, University of California, Santa Barbara, California, USA

¹⁶Arctic and Antarctic Research Institute, Saint Petersburg, Russia

¹⁷Mediterranean Institute for Advanced Studies, IMEDEA, CSIC-UIB, Mallorca, Spain

¹⁸Laboratoire Interuniversitaire des Systèmes Atmosphériques, Université de Paris7 et Université de Paris12, Creteil, France

¹⁹Laboratoire des Sciences du Climat et de l'Environnement, Gif-sur-Yvette, France

²⁰Physics Department, University of Miami, Coral Gables, Florida, USA

²¹Institute for Meteorology, University of Hamburg, Hamburg, Germany

²²NOAA/NESDIS Center for Satellite Applications and Research (STAR), Camp Springs, Maryland, USA

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



²³Department of Physics and Astronomy, Howard University, Washington, DC, USA

²⁴Space Physics Laboratory, Vikram Sarabhai Space Centre, Trivandrum, India

²⁵Department of Atmospheric Sciences, University of Washington, Seattle, Washington, USA

²⁶P. P. Shirshov Institute of Oceanology, Russian Academy of Sciences, Moscow, Russia

²⁷Space Physics Laboratory, Taras Shevchenko National University of Kyiv, Kyiv, Ukraine

²⁸Institut Maurice-Lamontagne, Mont-Joli, Quebec, Canada

²⁹Département de biologie, chimie et géographie, Université du Québec à Rimouski, Rimouski, Québec, Canada

³⁰Bedford Institute of Oceanography, Bedford, Nova Scotia, Canada

³¹Atmospheric Chemistry and Dynamics Branch, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

³²Climate and Radiation Branch, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

³³Marine Meteorology Division, Naval Research Laboratory, Monterey, California, USA

³⁴Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado, USA

³⁵University of North Dakota, Grand Forks, North Dakota, USA

³⁶Science Systems and Applications, Inc., Lanham, Maryland, USA

³⁷Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, USA

Received: 21 December 2010 – Accepted: 24 December 2010 – Published: 8 January 2011

Correspondence to: A. Smirnov (alexander.smirnov-1@nasa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

The Maritime Aerosol Network (MAN) has been collecting data over the oceans since November 2006. Over 80 cruises were completed through early 2010 with deployments continuing. Measurements areas included various parts of the Atlantic Ocean, the Northern and Southern Pacific Ocean, the South Indian Ocean, the Southern Ocean, the Arctic Ocean and inland seas. MAN deploys Microtops hand-held sunphotometers and utilizes a calibration procedure and data processing traceable to AERONET. Data collection included areas that previously had no aerosol optical depth (AOD) coverage at all, particularly vast areas of the Southern Ocean. The MAN data archive provides a valuable resource for aerosol studies in maritime environments. In the current paper we present results of AOD measurements over the oceans, and make a comparison with satellite AOD retrievals and model simulations.

1 Introduction

Atmospheric aerosol optical studies, involving radiative forcing analysis, aerosol-cloud interactions, remote sensing of the atmosphere, and global aerosol modeling require accurate information on aerosol optical depth (AOD). Sea salt is a major contributor to the columnar AOD over the oceans (Mahowald et al., 2006), and therefore affects the radiation budget directly (e.g., Haywood et al., 1999) and indirectly (O'Dowd et al., 1999). The complexity of aerosol production (Lewis and Schwartz, 2004) and advection from land sources warrant systematic measurements of aerosol optical parameters in maritime environments. Statistical robustness is required to better understand regional aerosol climatology and trends derived from the long-term satellite records.

Generally speaking, not all areas of the World Ocean can be studied from islands; aside from environmental satellites, ships are the only platform whereby measurements can be obtained. Ideally, a long-term comprehensive program is needed to include AOD on the list of routine meteorological and/or scientific measurements carried out

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al. (2003). The estimated uncertainty of the optical depth in each channel does not exceed ± 0.02 (Knobelspiesse et al., 2004), primarily due to inter-calibration against AERONET reference CIMEL instruments that are accurate to ~ 0.005 at most wavelengths (Eck et al., 1999). Microtops II instruments have shown good calibration stability over the years. Most of the instruments were manufactured in the late 1990s and have the original filters in place. The variability in calibration coefficients within a few percent over three years relative to AERONET reference CIMELs is quite acceptable. Figure 1 shows the variability in calibration coefficients (extraterrestrial irradiance signal, V_0) for one particular Microtops II. Certain changes in the calibration (post-field deployment in particular) are typically associated with aerosol deposition on the optics window that occurs at sea. After window cleaning, the calibration coefficients often approximate their original (pre-deployment) values. However, for some instruments we occasionally observed filter degradation which manifests itself as a rapid changes in the calibration coefficient.

The Maritime Aerosol Network measurement area has included northern and southern parts of the Atlantic Ocean; transects north–south, south–north, and east–west in the Pacific; intensive study areas in the Southern Ocean and off the coast of Antarctica including a number of circumnavigation cruises in high southern latitudes. A cruise area in the South Indian Ocean included the region between Reunion, Crozet, Kerguelen and Amsterdam Islands, as well as in the Mozambique Channel. Atmospheric measurements in the Bay of Bengal, Gulf of Bothnia as well as the Arabian, Mediterranean, Black, Baltic, Norway, Bering, Beaufort Seas, represented important contributions to the database. Previously, some of those oceanic regions (e.g. the Bering Sea, the Beaufort Sea, the South Indian Ocean, coast of Antarctica) had very limited or no surface-based AOD coverage at all.

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

accumulating data, with over 1700 days of measurements as of March 2010. The Level 2.0 data archive is mapped in Fig. 2.

Figure 3 shows latitudinal dependence of the AOD for different oceans and seas. The oceanic data have been divided roughly by longitude between three oceans: Atlantic (20° E–70° W), Pacific (70° W–150° E), and Indian (20° E–150° E). We consider measurements made south of 60° S as belonging to the Southern Ocean. Data collection in the Arctic Ocean was limited to the Beaufort Sea area. Measurements taken over Baltic, Black and Mediterranean Seas were grouped together. The measured aerosol properties (AOD, Angstrom parameter, and coarse mode fraction of AOD) for each area are shown in Figs. 3–6.

Figure 3a shows the latitudinal dependence of AOD daily averages at a wavelength 500 nm over the Atlantic Ocean. One can observe a pronounced peak in the distribution within the latitudinal belt 5°–20° N influenced by the dust and biomass burning aerosol transport from northern Africa. Optical depth variability is rather high ranging from typical values for the remote regions (~ 0.07 at 500 nm) to high aerosol loading close to 1.0. In the Southern Hemisphere $\tau_a(500)$ is typically less than 0.10; in some cases being as low as 0.04. Aerosol optical properties in the area north of 30° N are highly variable probably due to the various pollution aerosol sources in Europe and episodic dust transport from Africa. A few measurements in the northern areas near Greenland and Spitsbergen in the summer months yielded $\tau_a(500)$ values ~ 0.07 (typical of background conditions over the oceans). The $\tau_a(500)$ frequency histogram (bin size $\delta\tau_a = 0.05$) in Fig. 4a shows a peak at $\tau_a(500) \sim 0.075$ and indicates that 75% of the data has $\tau_a(500) < 0.20$. However the distribution has a “tail” that contains 25% of the daily averages. Marine and dust aerosol were clearly influential in producing the small value of the Angstrom parameter (~ 0.3) seen at the peak of the frequency distribution (bin size $\delta\alpha = 0.20$) of Fig. 5a. The spectral de-convolution algorithm allows the partition of aerosol optical depth into fine and coarse parts. The SDA-estimated coarse mode fraction (ratio of coarse mode AOD to total AOD at a wavelength of 500 nm) varies mainly within the 0.6–0.8 range in regions where marine and dust aerosols

**Maritime Aerosol
Network as
a component of
AERONET**A. Smirnov et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

are dominant while being significantly different for the regions where pollution and biomass burning aerosols to be more predominant (0.2–0.4). Frequency distribution of the coarse mode fraction (Fig. 6a) (bin size = 0.1) peaks at 0.75 with over 60% of occurrences within the 0.5–0.8 range.

According to the AERONET island stations in the Pacific and available publications summarized by Smirnov et al. (2002), the remote areas are typically characterized by $\tau_a(500)$ values ~ 0.07 . Overall this observation was confirmed by MAN ship-based measurements (Fig. 3b). However aerosol optical properties were highly variable (AODs ranged from 0.10 to 0.45) near Japan (heavily influenced by pollution aerosol and dust from Asia), and were occasionally elevated (a few days with AODs ~ 0.20 and higher were recorded) near the island of Hawaii, and the average AOD was ~ 0.20 near the west coast of South America at tropical latitudes. In the Pacific the bi-modal frequency distributions of $\tau_a(500)$ and the Angstrom parameter (Figs. 4b and 5b) are indicative of two optical conditions. The first is associated with the remote ocean with peak values near $\tau_a(500) \sim 0.07$ and $\alpha \sim 0.5$. The second modal feature has a most probable optical depth of ~ 0.23 and $\alpha \sim 1.0$. Therefore, in this latter case the fine mode aerosol fraction contributes more than 50% to the total AOD. The coarse mode fraction frequency distribution (Fig. 6b) peaks at 0.65 however it is wider than in the Atlantic (Fig. 6a) with almost equal frequencies within the range of 0.2–0.6.

Significant progress has been made in data collection over the Indian Ocean (Fig. 3c). Measurements over the Arabian Sea and the Bay of Bengal demonstrated a variety of optical conditions. AOD values were quite high with daily averages largely over 0.20. Overall the AOD over the Bay of Bengal was higher and also produced greater spectral dependence (high Angstrom parameter indicative of a dominant fine mode particle contribution). Optical properties in the region between Reunion and Tromelin islands and in the Mozambique Channel were highly variable. Continental aerosol plumes frequently appear in the area with at least one possible case of biomass burning aerosol from Africa (12 November 2009) with $\tau_a(500) \sim 0.60$ and $\alpha \sim 1.4$. Measurements in the South Indian Ocean in the area between Reunion,

**Maritime Aerosol
Network as
a component of
AERONET**A. Smirnov et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

not impossible to reproduce using other remote sensing techniques. Measurements in the Southern Ocean are comparable with the AERONET-based and other (see Tomasi et al., 2007) coastal measurements in Antarctica [$\tau_a(500)\sim 0.02\text{--}0.03$].

Several cruises conducted in the Mediterranean, Black, Baltic Seas including the Gulf of Bothnia provided a useful but relatively small dataset. Aerosol optical depth was highly variable (Fig. 3e) changing mainly within 0.10–0.40 range, except for the Gulf of Bothnia where $\tau_a(500)$ was less than 0.10. Data collection in the Beaufort Sea area (Fig. 3f, north of 65° N) enabled the characterization of background conditions during the summer of 2007 ($\tau_a(500)\sim 0.04$) as well as capturing Arctic haze events in the spring of 2008. A variety of optical conditions, such as biomass burning aerosol transported from Alaska were found during the summer of 2009. Data acquired in the Bering Sea are included in Fig. 3f in order to provide a basis of comparison with other MAN data. While limited to only five days of measurements it shows $\tau_a(500)$ to be $\sim 0.06\text{--}0.08$ which is comparable with the remote Pacific Ocean data but higher than Beaufort Sea background results by approximately of a factor of 1.5 to 2. Figures 4f and 5f permit a direct comparison of the AOD and Angstrom parameters with other regions. Measurements presented in Figs. 3–6e,f were carried out in different regions and in various seasons when optical conditions were determined by a mixture of maritime and continental aerosols. The scatter of the aerosol optical parameters is evident but α in the majority of cases is higher than 1.0 which is an indication of the significant contribution of fine particles to the attenuation in the atmospheric column (the coarse mode fraction of AOD varies mainly within 0.1–0.5 range). The most frequent AOD is ~ 0.12 for both subsets (Fig. 4e, f) however AOD frequencies are skewed towards higher AODs in Fig. 4f and towards smaller AODs in the case of the Bering and Beaufort Seas (Fig. 4f).

Overall statistics for oceanic areas (we did not include inland seas – Baltic, Black, and Mediterranean) is presented in Fig. 7. Despite the fact that vast areas still have limited or no coverage we can delineate some general characteristic features of aerosol optical properties over the oceans:

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- AOD at a wavelength 500 nm is less than 0.10 over oceanic areas not influenced by continental pollution, smoke or dust outflows – Fig. 7a.
- The Angstrom parameter (a general indicator of aerosol particle size) is generally smaller (<0.50) than values reported over continents (Holben et al., 2001) and in many instances less than values reported over island sites (Smirnov et al., 2002, 2009) – Fig. 7b. Desert dust contributed about 10% to the overall daily statistics if we consider $\text{AOD} > 0.2$ and $\alpha < 0.6$ as thresholds for a dust subset.
- The coarse mode AOD at 500 nm is less than 0.10 for vast majority of occurrences – Fig. 7c.
- Over 50% of the coarse mode fraction is within the range 0.50–0.80, denoting dominance of coarse aerosol in the total aerosol optical depth – Fig. 7d.

4 Comparison with satellite retrievals and global transport models

The ship-borne measurements provide an excellent opportunity for comparison with global aerosol transport models and satellite retrievals. AOD differences between satellite retrievals or model simulations and ship-borne AODs are presented in this section. In order to better visualize comparisons we present AOD differences as a function of latitude against MAN ground-truth for each sensor or model. Sunphotometer measurement series (Level 2.0) were spectrally adjusted using log-linear interpolation to the “validation” wavelength of 550 nm.

The global model GOCART is driven by the assimilated meteorological fields from the Goddard Earth Observing System Data Assimilation System (GEOS4-DAS) and simulates major aerosol types of dust, sulfate, black carbon, organic matter, and sea salt (details described in Chin et al., 2002, 2009, and references therein). GOCART simulated aerosol optical depth used in this study is archived at 1° latitude by 1.25° longitude spatial resolution every three hours. For comparisons in this study, the GOCART output was extracted to match the MAN observations at the closest location and time.

**Maritime Aerosol
Network as
a component of
AERONET**A. Smirnov et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

GEOS-Chem (www.geos-chem.org) is a global chemical transport model driven by assimilated meteorology from the NASA Global Modeling and Assimilation Office (GMAO). Simulations shown here were performed with v8-03-01 of the model with GEOS-5 meteorology at $2^\circ \times 2.5^\circ$ horizontal resolution (degraded from $0.5^\circ \times 0.67^\circ$) and 47 vertical levels. The total AOD shown here includes contributions from sulfate, nitrate, ammonium, black carbon, organic carbon, sea salt and dust. Aerosol optical properties are based on the Global Aerosol Data Set (GADS) (Kopke et al., 1997) with modifications from Drury et al. (2010) and Jaegle et al. (2010). Model output is sampled along the MAN ship tracks and matched temporally within 30 min (the chemistry time step of the model).

The comparison with the median AeroCom model (Schulz et al., 2006) constructed from the output representing year 2000 simulations by twelve models (GISS, GOCART, KYU, LOA, LSCE, MATCH, MOZGN, MPI.HAM, PNNL, TM5_B, UIO_CTM, UMI; see details in Textor et al., 2006; Kinne et al., 2006) was made in the following way. MAN data from a given day were averaged per day and the mean latitude/longitude position was calculated. AeroCom median model data were extracted for a corresponding month when ship-based observations were made and at the mean MAN location for any given day with observations available. Each day with a MAN observation thus has one corresponding model value in a $1^\circ \times 1^\circ$ grid. This “matching” was thus done differently from other models and satellite sensors but respects seasonal variability.

The number of morning (Terra) and afternoon (Aqua) MODIS retrievals matching ship-based τ_a was quite high. The matchup criteria were a modification of Ichoku et al. (2004). We looked for any series of MAN measurements within ± 30 min of the MODIS overpass time. MODIS was required to retrieve at least 5 out of 25 pixels in the 50 km box around the ship location (details are presented by Kleidman et al., 2010). In the case of multiple matching sunphotometer measurement series we took the one closest in time to the overpass if the AOD variability was small and averaged MAN series measurements if variability was large after eliminating outliers. In over 90% of the cases we selected the closest series.

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The matchup criteria for the MISR (algorithm version 22) product included successful retrievals either in the 17.6 km MISR retrieval region containing selected ship-based measurement (the “central” region), or in one or more of the eight retrieval regions surrounding the central one. The MAN time series for each coincidence include at least one AOD measurement during the hour before the MISR overpass, and at least one during the hour after the overpass (Kahn et al., 2005, 2010). The number of matching cases for MISR is limited (only 61 match-ups) with several outliers. Five out of seven outliers were identified as being cloud contaminated (Kahn et al., 2010). The proximity of a coastline in one case and an ice surface in the other case complicated the retrieval process for the other two outliers.

Zhang and Reid (2006) developed a methodology to minimize cloud contamination and other biases in MODIS aerosol product for implementation in operational aerosol data assimilation (DA). This DA quality level-3 Terra MODIS and Aqua MODIS AODs (Zhang and Reid, 2006; Shi et al., 2010) will be used in this study (marked as DA – data assimilation quality assured). The over ocean collection 5 MODIS level-2 AODs (marked as Standard) are included for comparison. We consider any pairs of MODIS and MAN series data within ± 30 min of the overpass time and spatially within 30 km. If more than one MAN series data point is available then we pick the closest in time.

The temporal and spatial scale differences between point measurements from MAN and area-grids from satellite retrievals and model simulations may lead to some differences. The temporal difference is addressed by utilizing the MAN series data within a set period from the satellite or model reporting time. However, the spatial difference can only be addressed with several widely distributed measurements within the domain. As a result, some of the measurements from MAN may capture episodic aerosol plumes, which may not be detected by larger grid scale products that average over a large region. The spatial difference would tend to affect periods when MAN reported higher AOD and difference would be greater for the largest area-grids.

Figure 8 presents AOD differences between global model simulations and ship-borne AODs as a function of latitude. GOCART and GEOS-Chem simulations were available

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



only for the year 2007. From Fig. 8a one can observe that GOCART overestimates AOD more often. Positive bias is evident south of 60° S and north of 30° N. A significant temporal variability in areas of the Atlantic influenced by dust and biomass burning sources produced almost symmetrical AOD differences within the belt 0° – 30° N. While the GEOS-Chem model (Fig. 8b) reproduces many of the NH observations (clustering around zero bias), large negative excursions are also evident in plumes measured in 2007. The complexity of various continental sources, as well as the coarser horizontal resolution of the model simulation might explain the disparity. Unlike GOCART the GEOS-Chem and sunphotometer AOD differences are approximately equally distributed around zero south of 60° S. The median AEROCOM model shows more scatter mainly in the Southern Hemisphere (Fig. 8c). AOD differences are mostly negative, however distributed almost equally in the areas north of the equator to 30° N.

Satellite retrievals from MODIS and MISR (Fig. 9a–b) indicate a positive bias, very similar to each other, although the MODIS differences are smaller. Separating Terra and Aqua retrievals would show that Terra is more severely biased high than Aqua (Remer et al., 2008). MISR and MODIS retrievals are more likely to be biased high than low over ocean, as the algorithms assume cloud-free scene and dark surfaces, whereas unscreened cloud or whitecaps, and non-zero surface reflectivity due to runoff, pollution, or biological activity, would all increase scene reflectance (Kahn et al., 2007).

The standard and data assimilation quality product for Terra MODIS comparison (Fig. 9c–d) shows a significant improvement in the latitudinal dependence of AOD differences. The noticeable positive bias in the Standard AOD (Fig. 9c) disappeared in the DA product (Fig. 9d). In the Northern Hemisphere differences are almost evenly distributed around zero with the 0 – 60° degrees belt, while changing the sign of the AOD differences further north of 60° N. The strong positive bias in the Southern Ocean for the Standard AOD became much smaller for the DA product. Comparison made for the Aqua MODIS (Fig. 9e–f) does show some improvements but no drastic changes. A number of outliers on the negative side might be associated with the unnoticed cloud contamination of the sunphotometer data.

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We would like to emphasize that our analysis is not intended to determine how many retrievals are within the claimed uncertainty boundaries or beyond. Rather we wanted to show where satellite retrieval biases exist and, in what latitudinal belts corrections are needed. For example, in the southern latitudes (south of 40 degrees) the sunphotometer AODs are low compared with satellite retrievals and modeling results. This discrepancy can, at least partly, be explained by uncertainties in aerosol production rates (Lewis and Schwartz, 2004), foam formation and its latitudinal distribution (Anguelova and Webster, 2006), by a process of quality control that excludes some residual cloud contamination (Zhang and Reid, 2010), by the accuracy of radiative transfer models used (Melin et al., 2010), and more accurate accounting for surface reflectance effects (Sayer et al., 2010).

A valid comparison among various models, satellite products and sunphotometer measurements (SP) is presented in Fig. 10. The frequency of occurrences histogram indicates that vast majority of the differences are positive. Only two out of nine differences (DA Terra – SP and AEROCOM – SP) are biased slightly negative. The AEROCOM – SP difference has a much wider distribution and as a consequence peaks at only 20%, lower than others. GEOS-Chem is almost symmetrical around zero, although biased slightly high as are the other models and sensors. The MISR – SP distribution shows bi-modality mainly because of the small number of match-ups available.

5 Summary

The Maritime Aerosol Network has continued extended spectral AOD data collection to areas that previously had no coverage. A web-based data archive provides the international scientific community with valuable data for satellite retrieval validation, atmospheric correction and other applications. Many areas of the World Ocean still have little or no coverage and our objective in the future is to extend coverage to all of these regions. Our international, multi-institutional collaborative effort will significantly

enhance our knowledge on the global aerosol distribution over the oceans. We foresee a continuation of this effort on various ships of opportunity.

Acknowledgement. The authors thank Hal Maring (NASA Headquarters) for his support of AERONET. The authors would like to acknowledge managerial and operational support from M. Sorokin, A. Scully, A. Tran, P. Kenny, D. Hamilton, L. Bariteau, R. Dunn, M. Conley, P. Schoessow, H. Gomes, L. Logan, M. Reynolds, A. Flores, D. A. Siegel, A. Proshutinsky, L. Rainville, A. Jayakumar, S. Schick, D. Menzies, E. Emry, C. Swan, K. G. Fairbarn (USA); M. Panchenko, O. Kopelevich, A. Sinitsyn, D. Kabanov, A. Tikhomirov, A. Kalsin, S. Terpu-
gova, V. Polkin (Sr), V. Polkin (Jr), N. Vlasov, Yu. Turchinovich, A. Gubin, Y. Zyulyaeva (Russia);
P. Goloub, L. Blarel, S. Triquet, P. Hernandez, V. Dufлот, T. Lecointre, S. Barataud, P. Ricaud,
P. Sangiardi, A. Kartavtseff, J.-F. Ternon, F. Jourdin, C. Petus, J. Nicolas, S. Devidal, L. Marti-
non, M. Faillot, F. Gabarrot, N. Villeneuve, I. Jubert, M. Barblu, G. Duval (France); C. Powell,
C. Gallienne (UK); C. Schlosser, Y. Zoll, M. Schlundt, M. Heller, T. Hanschmann, K. Lengfeld,
A. Tessorodorf, N. Renkosik, T. Heus, K. Lonitz, B. Quack, T. Dinter, A. Wassmann, M. Schlundt,
B. Pospichal, F. Wittrock (Germany); A. Bromley, R. Martin, G. Brailsford (New Zealand);
J. Kowalczyk, A. Ponczkowska, J. Pasnicki, K. Zielinski, P. Makuch, B. Lednicka (Poland);
S. Babu, S. K. Satheesh, V. S. Nair, S. N. Beegum (India), S. Piketh, D. Williams, B. Kuyper,
E. Robertson (South Africa), L. Jankowski, R. Matarrese (Italy), R. M. Castillo (Spain). One of
the co-authors (Jean Sciare) would like to thank Institut Polaire Francais (IPEV) for the support
provided within the AEROTRACE project.

References

- Anguelova, M. D. and Webster, F.: Whitecap coverage from satellite measurements: A first step toward modeling the variability of oceanic whitecaps, *J. Geophys. Res.*, 111, C03017, doi:10.1029/2005JC003158, 2006.
- Barteneva, O. D., Nikitinskaya, N. I., Sakunov, G. G., and Veselova, L. K.: Atmospheric transmittance in the visible and near IR spectral range, *Gidrometeoizdat, Leningrad*, 224 pp., 1991 (in Russian).
- Chin, M., Ginoux, P., Kinne, S., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: Tropospheric aerosol optical thickness from the GOCART

AMTD

4, 1–32, 2011

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

model and comparisons with satellite and sunphotometer measurements, *J. Atmos. Sci.*, 59, 461–483, 2002.

Chin, M., Diehl, T., Dubovik, O., Eck, T. F., Holben, B. N., Sinyuk, A., and Streets, D. G.: Light absorption by pollution, dust, and biomass burning aerosols: a global model study and evaluation with AERONET measurements, *Ann. Geophys.*, 27, 3439–3464, doi:10.5194/angeo-27-3439-2009, 2009.

Drury, E., Jacob, D. J., Spurr, R. J. D., Wang, J., Shinozuka, Y., Anderson, B. E., Clarke, A. D., Dibb, J., McNaughton, C., and Weber, R.: Synthesis of satellite (MODIS), aircraft (ICARTT), and surface (IMPROVE, EPA-AQS, AERONET) aerosol observations over Eastern North America to improve MODIS aerosol retrievals and constrain aerosol concentrations and sources, *J. Geophys. Res.*, 115, D14204, doi:10.1029/2009JD012629, 2010.

Eck, T. F., Holben, B. N., Reid, J. S., Dubovik, O., Smirnov, A., O'Neill, N. T., Slutsker, I., and Kinne, S.: Wavelength dependence of the optical depth of biomass burning, urban, and desert dust aerosol, *J. Geophys. Res.*, 104, 31333–31350, 1999.

Haywood, J., Ramaswamy, V., and Soden, B.: Tropospheric aerosol climate forcing in clear-sky satellite observation over the oceans, *Science*, 283, 1299–1303, 1999.

Holben, B. N., Eck, T. F., Slutsker, I., Tanre, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y., Nakajima, T., Lavenue, F., Jankowiak, I., and Smirnov, A.: AERONET – a federated instrument network and data archive for aerosol characterization, *Remote. Sens. Environ.*, 66, 1–16, 1998.

Holben, B. N., Tanre, D., Smirnov, A., Eck, T. F., Slutsker, I., Abuhassan, N., Newcomb, W. W., Schafer, J., Chatenet, B., Lavenue, F., Kaufman, Y. J., Van de Castle, J., Setzer, A., Markham, B., Clark, D., Frouin, R., Halthore, R., Karnieli, A., O'Neill, N. T., Pietras, C., Pinker, R. T., Voss, K., and Zibordi, Z.: An emerging ground-based aerosol climatology: aerosol optical depth from AERONET, *J. Geophys. Res.*, 106, 12067–12097, 2001.

Ichoku, C., Kaufman, Y. J., Remer, L. A., and Levy, R.: Global aerosol remote sensing from MODIS, *Adv. Space Res.*, 34, 820–827, 2004.

Jaeglé, L., Quinn, P. K., Bates, T., Alexander, B., and Lin, J.-T.: Global distribution of sea salt aerosols: new constraints from in situ and remote sensing observations, *Atmos. Chem. Phys. Discuss.*, 10, 25687–25742, doi:10.5194/acpd-10-25687-2010, 2010.

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kahn, R. A., Gaitley, B., Martonchik, J., Diner, D., Crean, K., and Holben, B.: MISR global aerosol optical depth validation based on two years of coincident AERONET observations, *J. Geophys. Res.*, 110, D10S04, doi:10.1029/2004JD004706, 2005.
- 5 Kahn, R. A., Garay, M. J., Nelson, D. L., Yau, K. K., Bull, M. A., Gaitley, B. J., Martonchik, J. V., and Levy, R. C.: Satellite-derived aerosol optical depth over dark water from MISR and MODIS: comparisons with AERONET and implications for climatological studies, *J. Geophys. Res.*, 112, D18205, doi:10.1029/2006JD008175, 2007.
- 10 Kahn, R. A., Gaitley, B. J., Garay, M., Diner, D. J., Eck, T. F., Smirnov, A., and Holben, B. N.: Multi-angle Imaging SpectroRadiometer global aerosol product assessment by comparison with the Aerosol Robotic Network, *J. Geophys. Res.*, 115, D23209, doi:10.1029/2010JD014601, 2010.
- 15 Kinne, S., Schulz, M., Textor, C., Guibert, S., Balkanski, Y., Bauer, S. E., Bernsten, T., Berglen, T. F., Boucher, O., Chin, M., Collins, W., Dentener, F., Diehl, T., Easter, R., Feichter, J., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Herzog, M., Horowitz, L., Isaksen, I., Iversen, T., Kirkevåg, A., Kloster, S., Koch, D., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Lesins, G., Liu, X., Lohmann, U., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T., and Tie, X.: An AeroCom initial assessment – optical properties in aerosol component modules of global models, *Atmos. Chem. Phys.*, 6, 1815–1834, doi:10.5194/acp-6-1815-2006, 2006.
- 20 Kleidman, R. G., Smirnov, A., Levy, R. C., Mattoo, S., and Tanre, D.: Evaluation and wind speed dependence of MODIS aerosol retrievals over open ocean, *IEEE Trans. Geosci. Remote Sens.*, submitted, 2010.
- Knobelspiesse, K. D., Pietras, C., and Fargion, F. S.: Sun-pointing error correction for sea deployment of the Microtops II handheld Sun photometer, *J. Atmos. Oceanic Technol.*, 20, 767–771, 2003.
- 25 Knobelspiesse, K. D., Pietras, C., Fargion, G. S., Wang, M. H., Frouin, R., Miller, M. A., Subramaniam, S., and Balch, W. M.: Maritime aerosol optical thickness measured by handheld sunphotometers, *Remote Sens. Environ.*, 93, 87–106, 2004.
- Kopke, P., Hess, M., Schult, I., and Shettle, E. P.: Global aerosol data set, Max Planck Inst. fur Meteorol., Hamburg, Germany, 44 pp., 1997.
- 30 Lewis, E. R. and Schwartz, S. E.: Sea salt aerosol production: mechanisms, methods, measurements, and models – a critical review, American Geophysical Union, Washington, DC, 413 pp., 2004.

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mahowald, N. M., Lamarque, J.-F., Tie, X. X., and Wolff, E.: Sea-salt aerosol response to climate change: Last Glacial Maximum, pre-industrial and doubled carbon dioxide climates, *J. Geophys. Res.*, 111, D05303, doi:10.1029/2005JD006459, 2006.

Melin, F., Clerici, M., Zibordi, G., Holben, B. N., and Smirnov, A.: Validation of SeaWiFS and MODIS aerosol products with globally distributed AERONET data, *Remote Sens. Environ.*, 114, 230–250, 2010.

Morys, M., Mims, F. M., Hagerup, S., Anderson, S. E., Baker, A., Kia, J., and Walkup, T.: Design, calibration, and performance of MICROTUPS II handheld ozone monitor and Sun photometer, *J. Geophys. Res.*, 106, 14573–14582, 2001.

O'Dowd, C. D., Lowe, J. A., and Smith, M. H.: Coupling sea-salt and sulphate interactions and its impact on cloud droplet concentration predications, *Geophys. Res. Lett.*, 26, 1311–1314, doi:10.1029/1999GL900231, 1999.

O'Neill, N. T., Eck, T. F., Holben, B. N., Smirnov, A., Dubovik, O., and Royer, A.: Bimodal size distribution influences on the variation of Angstrom derivatives in spectral and optical depth space, *J. Geophys. Res.*, 106, 9787–9806, 2001.

O'Neill, N. T., Eck, T. F., Smirnov, A., Holben, B. N., and Thulasiraman, S.: Spectral discrimination of coarse and fine mode optical depth, *J. Geophys. Res.*, 108(D17), 4559, doi:10.1029/2002JD002975, 2003.

Porter, J. N., Miller, M., Pietras, C., and Motell, G.: Ship-based sunphotometer measurements using Microtops sunphotometers, *J. Atmos. Oceanic Technol.*, 18, 765–774, 2001.

Remer, L. A., Kleidman, R. G., Levy, R. C., Kaufman, Y. J., Tanre, D., Matoo, S., Martins, J. V., Ichoku, C., Koren, I., Yu, H. B., Holben, B. N.: Global aerosol climatology from the MODIS satellite sensors, *J. Geophys. Res.*, 113, D14S07, doi:10.1029/2007JD009661, 2008.

Sayer, A. M., Thomas, G. E., and Grainger, R. G.: A sea surface reflectance model for (A)ATSR, and application to aerosol retrievals, *Atmos. Meas. Tech.*, 3, 813–838, doi:10.5194/amt-3-813-2010, 2010.

Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S., Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, S., Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-industrial simulations, *Atmos. Chem. Phys.*, 6, 5225–5246, doi:10.5194/acp-6-5225-2006, 2006.

Shi, Y., Zhang, J., Reid, J. S., Holben, B., Hyer, E. J., and Curtis, C.: An analysis of the Collection 5 MODIS over-ocean aerosol optical depth product for its implication in aerosol

assimilation, *Atmos. Chem. Phys. Discuss.*, 10, 20239–20265, doi:10.5194/acpd-10-20239-2010, 2010.

Smirnov, A., Holben, B. N., Kaufman, Y. J., Dubovik, O., Eck, T. F., Slutsker, I., Pietras, C., and Halthore, R.: Optical properties of atmospheric aerosol in maritime environments, *J. Atmos. Sci.*, 59, 501–523, 2002.

Smirnov, A., Holben, B. N., Lyapustin, A., Slutsker, I., and Eck, T. F.: AERONET processing algorithm refinement, AERONET Workshop, El Arenosillo, Spain, 10–14 May 2004.

Smirnov, A., Holben, B. N., Sakerin, S. M., Kabanov, D. M., Slutsker, I., Chin, M., Diehl, T. L., Remer, L. A., Kahn, R. A., Ignatov, A., Liu, L., Mishchenko, M., Eck, T. F., Kucsera, T. L., Giles, D. M., and Kopelevich, O. V.: Ship-based aerosol optical depth measurements in the Atlantic Ocean, comparison with satellite retrievals and GOCART model, *Geophys. Res. Lett.*, 33, L14817, doi:10.1029/2006GL026051, 2006.

Smirnov, A., Holben, B. N., Slutsker, I., Giles, D. M., McClain, C. R., Eck, T. F., Sakerin, S. M., Macke, A., Croot, P., Zibordi, G., Quinn, P. K., Sciare, J., Kinne, S., Harvey, M., Smyth, T. J., Piketh, S., Zielinski, T., Proshutinsky, A., Goes, J. I., Nelson, N. B., Larouche, P., Radionov, V. F., Goloub, P., Moorthy, K. K., Matarrese, R., Robertson, E. J., and Jourdin, F.: Maritime Aerosol Network as a component of Aerosol Robotic Network, *J. Geophys. Res.*, 114, D06204, doi:10.1029/2008JD011257, 2009.

Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Bernsten, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, I., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles within AeroCom, *Atmos. Chem. Phys.*, 6, 1777–1813, doi:10.5194/acp-6-1777-2006, 2006.

Tomasi, C., Vitale, V., Lupi, A., Di Carmine, C., Campanelli, M., Herber, A., Treffeisen, R., Stone, R. S., Andrews, E., Sharma, S., Radionov, V., von Hoyningen-Huene, W., Stebel, K., Hansen, G. H., Myhre, C. L., Wehrli, C., Aaltonen, V., Lihavainen, H., Virkkula, A., Hillamo, R., Strom, J., Toledano, C., Cachorro, V. E., Ortiz, P., de Frutos, A. M., Blindheim, S., Frioud, M., Gausa, M., Zielinski, T., Petelski, T., and Yamanouchi, T.: Aerosols in polar regions: a historical overview based on optical depth and in situ observations, *J. Geophys. Res.*, 112, D16205, doi:10.1029/2007JD008432, 2007.

AMTD

4, 1–32, 2011

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Vinoj, V., Anjan, A., Sudhakar, M., Satheesh, S. K., Srinivasan, J., and Moorthy, K. K.: Latitudinal variation of aerosol optical depths from northern Arabian Sea to Antarctica, *Geophys. Res. Lett.*, 34, L10807, doi:10.1029/2007GL029419, 2007.

5 Zhang, J. and Reid, J. S.: MODIS Aerosol product analysis for data assimilation: assessment of Level 2 aerosol optical thickness retrievals, *J. Geophys. Res.*, 111, D22207, doi:10.1029/2005JD006898, 2006.

Zhang, J. and Reid, J. S.: A decadal regional and global trend analysis of the aerosol optical depth using a data-assimilation grade over-water MODIS and Level 2 MISR aerosol products, *Atmos. Chem. Phys.*, 10, 10949–10963, doi:10.5194/acp-10-10949-2010, 2010.

AMTD

4, 1–32, 2011

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. Microtops (S/N 3657) calibration history.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

⏴ ⏵

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.




Fig. 2. Maritime Aerosol Network global coverage – cruise tracks and daily averages of aerosol optical depth at 500 nm (squares are colored with respect to AOD values, i.e. blue – $AOD < 0.10$, green – $0.1 \leq AOD < 0.2$, yellow – $0.2 \leq AOD < 0.3$, orange – $0.3 \leq AOD < 0.5$, red – $0.5 \leq AOD < 0.7$, purple – $AOD \geq 0.7$).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 3. Latitudinal dependence of daily averaged aerosol optical depth in the Atlantic Ocean **(a)**, Pacific Ocean **(b)**, Indian Ocean **(c)**, Southern Ocean **(d)**, Baltic, Black, and Mediterranean Seas **(e)**, Bering and Beaufort Seas **(f)**.

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.




Fig. 4. Frequency of occurrences of daily averaged aerosol optical depth at 500 nm for **(a)** Atlantic Ocean, **(b)** Pacific Ocean, **(c)** Indian Ocean, **(d)** Southern Ocean, **(e)** Baltic, Black, and Mediterranean Seas, **(f)** Bering and Beaufort Seas.




Fig. 5. Frequency of occurrences of daily averaged Angstrom parameter for **(a)** Atlantic Ocean, **(b)** Pacific Ocean, **(c)** Indian Ocean, **(d)** Southern Ocean, **(e)** Baltic, Black, and Mediterranean Seas, **(f)** Bering and Beaufort Seas.




Fig. 6. Frequency of occurrences of the daily averaged coarse mode fraction for **(a)** Atlantic Ocean, **(b)** Pacific Ocean, **(c)** Indian Ocean, **(d)** Southern Ocean, **(e)** Baltic, Black, and Mediterranean Seas, **(f)** Bering and Beaufort Seas.




Fig. 7. Frequency of occurrences of daily averaged AOD (a), Angstrom parameter (b), coarse mode AOD (c), coarse mode fraction of AOD (d).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.




Fig. 8. Latitudinal dependence of aerosol optical depth differences between various global aerosol transport models and sunphotometer.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 9. Latitudinal dependence of aerosol optical depth differences between various satellite sensors and sunphotometer.

Maritime Aerosol Network as a component of AERONET

A. Smirnov et al.




Fig. 10. Frequency of occurrences of aerosol optical depth differences between various models/sensors and sunphotometer.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion